

CLINICAL POLICY

Galcanezumab-gnlm

Clinical Policy: Galcanezumab-gnlm (Emgality)

Reference Number: MDN.CP.PHAR.404

Effective Date: 12.1.23

Last Review Date: 10.16.23

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Galcanezumab-gnlm (Emgality[®]) is a calcitonin gene-related peptide (CGRP) receptor antagonist.

FDA Approved Indication(s)

Emgality is indicated in adults for the:

- Preventive treatment of migraine
- Treatment of episodic cluster headache

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Emgality is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Migraine Prophylaxis (must meet all):

1. Diagnosis of episodic or chronic migraine;
2. Age \geq 18 years;
3. Failure of at least 2 of the following oral migraine preventative therapies, each for 8 weeks and from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine);
4. Emgality is not prescribed concurrently with Botox[®] or other injectable or oral CGRP inhibitors (e.g., Aimovig, Ajoovy[®], Vyepti[™], Nurtec[®], Ubrelvy[™]);
5. Dose does not exceed:
 - a. Loading dose: 240 mg (2 injections) once;
 - b. Maintenance dose: 120 mg (1 injection) once monthly.
6. If currently receiving treatment with Botox[®] for migraine prophylaxis and request is for concurrent use of Botox and Emgality (i.e., not switching from one agent to another), all of the following (a, b, and c):

- a. Sufficient evidence is provided from at least two high-quality*, published studies in reputable peer-reviewed journals or evidence-based clinical practice guidelines that provide all of the following (i – iv):
**Case studies or chart reviews are not considered high-quality evidence*
 - i. Adequate representation of the member’s clinical characteristics, age, and diagnosis;
 - ii. Adequate representation of the prescribed drug regimen;
 - iii. Clinically meaningful outcomes such as a reduction in monthly migraine or headache days;
 - iv. Appropriate experimental design and method to address research questions (see Appendix E for additional information);
- b. Member has experienced and maintained positive response to Botox monotherapy as evidenced by a $\geq 30\%$ reduction in migraine days per month from baseline following at least 2 quarterly injection (6 months) of Botox monotherapy;
- c. Despite Botox monotherapy, member continues to experience ≥ 4 migraine days per month and/or severe migraine headaches that result in disability and functional impairment;

Approval duration: 3 months

B. Episodic Cluster Headaches (must meet all):

1. Diagnosis of episodic cluster headaches
2. Provider’s attestation that member has a history of ≥ 2 cluster periods lasting from 7 days to 1 year each and separated by ≥ 3 months;
3. Age ≥ 18 years;
4. Failure of verapamil at a dose of 360 mg per day, unless contraindicated or clinically significant adverse effects are experienced;
5. Emgality is not prescribed concurrently with other injectable or oral CGRP inhibitors (e.g., Aimovig, Ajovy, Vyepti, Nurtec, Ubrelvy);
6. Dose does not exceed 300 mg (3 injections) once monthly.

Approval duration: 3 months

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND

criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Migraine Prophylaxis (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B);
2. Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline;
3. Emgality is not prescribed concurrently with Botox or other injectable or oral CGRP inhibitors (e.g., Aimovig, Ajoovy, Vyepiti, Nurtec, Ubrelvy);
4. If request is for a dose increase, new dose does not exceed 120 mg (1 injection) once monthly.

Approval duration: 6 months

B. Episodic Cluster Headaches (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B);
2. Member is responding positively to therapy as evidenced by a reduction in cluster headache attack frequency;
3. Member meets one of the following (a or b):
 - a. Member has not received more than 12 months of consecutive treatment;
 - b. It has been at least 3 months since the member last received Emgality;
4. Emgality is not prescribed concurrently with other injectable or oral CGRP inhibitors (e.g., Aimovig, Ajoovy, Vyepiti, Nurtec, Qulipta, Ubrelvy);
5. If request is for a dose increase, new dose does not exceed 300 mg (3 injections) once monthly.

Approval duration: 6 months (up to a total of 12 months per cluster period)

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):

- a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.PMN.53 for Medicaid

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 for Medicaid or evidence of coverage documents;
- B. Chronic cluster headaches.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CGRP: calcitonin gene-related peptide

FDA: Food and Drug Administration

ICHD: International Classification of Headache Disorder

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Anticonvulsants such as: divalproex (Depakote [®]), topiramate (Topamax [®]), valproate sodium	Migraine Prophylaxis <i>Refer to prescribing information or Micromedex</i>	<i>Refer to prescribing information or Micromedex</i>
Beta-blockers such as: propranolol (Inderal [®]), metoprolol (Lopressor [®])*, timolol, atenolol (Tenormin [®])*, nadolol (Corgard [®])*	Migraine Prophylaxis <i>Refer to prescribing information or Micromedex</i>	<i>Refer to prescribing information or Micromedex</i>
Antidepressants/tricyclic antidepressants* such as: amitriptyline (Elavil [®]), venlafaxine (Effexor [®])	Migraine Prophylaxis <i>Refer to prescribing information or Micromedex</i>	<i>Refer to prescribing information or Micromedex</i>
verapamil*	Episodic Cluster Headache 120 mg PO TID	360 mg/day

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Aimovig (erenumab-aaoe)	<p>Migraine Prophylaxis 70 mg SC once monthly</p> <p>Some patients may benefit from a dosage of 140 mg injected subcutaneously once monthly</p>	140 mg/month

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

*Off-label use

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity
- Boxed warning(s): none reported

Appendix D: General Information

- In clinical trials, a migraine day was defined as any calendar day in which the patient reported either a headache that lasted at least 2 consecutive hours and met International Classification of Headache Disorder (ICHD)-3 diagnostic criteria for migraine (with or without aura) or probable migraine (subtype in which only 1 migraine criterion is absent), or a day when a headache of any duration was treated with migraine-specific medications (triptans or ergots).
- Although Emgality given as either 120 mg SC once monthly or 240 mg SC once monthly showed a statistically significant decrease in migraine days per month compared to placebo as the primary outcome in the EVOLVE-1, EVOLVE-2, and REGAIN pivotal trials, there was no clinically significant difference between the two dosing regimens, and thus no significant additional benefit conferred from using a higher dose of Emgality. This is consistent with the FDA-approved maintenance dose of 120 mg SC once monthly.
- According to the ICHD-3 diagnostic criteria for cluster headaches, episodic cluster headaches occur in periods lasting from seven days to one year and are separated by periods of remissions that are at least 3 months. Chronic cluster headaches (affecting 10-15% of patients), on the other hand, occur for longer than a year without remission or with a remission that lasts less than 3 months. Of note, Emgality has only demonstrated efficacy in episodic cluster headaches. It failed to meet its primary endpoint in its chronic cluster headache phase 3 trial.

Appendix E: Appropriate Experimental Design Methods

- Randomized, prospective controlled trials are generally considered the gold standard; however:
 - In some clinical studies, it may be unnecessary or not feasible to use randomization, double-blind trials, placebos, or crossover.
 - Non-randomized prospective clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.

- Case reports and chart reviews are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Migraine prophylaxis	Loading dose: 240 mg SC once Maintenance dose: 120 mg SC once monthly	120 mg/month
Episodic cluster headaches	300 mg (administered as three consecutive injections of 100 mg each) SC at the onset of the cluster period, and then monthly until the end of the cluster period	300 mg/month

VI. Product Availability

- Single-dose prefilled pen: 120 mg/mL
- Single-dose prefilled syringe: 100 mg/mL, 120 mg/mL

VII. References

1. Emgality Prescribing Information. Indianapolis, IN: Eli Lilly and Company; Mar 2021. Available at: <http://www.emgality.com>. Accessed April 28, 2022.
2. Silberstein SD, Holland S, Freitag F, et al. American Academy of Neurology: Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults. *Neurology* 2012; 78: 1337-45.
3. Stauffer VL, Dodick DW, Zhang Q, et al. Evaluation of galcanezumab for the prevention of episodic migraine: the EVOLVE-1 randomized clinical trial. *JAMA Neurol.* 2018; 75(9):1080-1088.
4. Skljarevski V, Matharu M, Millen BA, et al. Efficacy and safety of galcanezumab for the prevention of episodic migraine: results of the EVOLVE-2 phase 3 randomized controlled clinical trial. *Cephalalgia.* 2018; 38(8):1442-1454.
5. Detke H, Wang S, Skljarevski V, et al. A phase 3, randomized, double-blind, placebo-controlled study of LY2951742 in patients with chronic migraine – the REGAIN study. Poster session presented at: International Headache Congress; Sept 7-10, 2017; Vancouver, Canada.
6. Headache Classification Committee of the International Headache Society. The International classification of headache disorders, 3rd edition (beta version). *Cephalalgia.* 2013; 33(9): 629-808.
7. Francis BJ, Becker WJ, and Pringsheim TM. Acute and preventative pharmacologic treatment of cluster headache. *Neurology.* 2010; 75: 463-473.
8. Robbins MS, Starling AJ, Pringsheim TM, Becker WJ, and Schwedt TJ. Treatment of cluster headache: The American Headache Society evidence-based guidelines. *Headache.* 2016; 56: 1093-1106.
9. Digre KB. The American Headache Society Position Statement On Integrating New Migraine Treatments Into Clinical Practice. *Headache* 2019; 59: 1-18.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3490	Unclassified drugs

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created from IL.PHAR.404 to align with HFS PDL.	10.16.23	

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.